

Changes in the refraction and visual acuity of patients during a period of the first month after phacoemulsification surgery

Dr. Pateras E.¹, Dr. Alonistiotis D.², Hyska B.³, Tsiantakou G.⁴

1. Associate Professor, Biomedical Department, Course of Optics and Optometry, University of West Attica UNIWA Athens
2. Ophthalmologist, Lecturer, 2nd University Clinic of Ophthalmology, "ATTIKON" Athens University General Clinic
3. Student, Biomedical Department, Course of Optics and Optometry, University of West Attica UNIWA Athens
4. Student, Biomedical Department, Course of Optics and Optometry, University of West Attica UNIWA Athens

Introduction

The crystalline lens of the eye is a transparent and biconvex structure, which is placed at the posterior chamber of the eye, between the iris and the vitreous humor^[1,2,3]. The lens is suspended in place by the radially arranged zonules fibers. Also, is anatomically composed by a surrounding capsule, the epithelial cells, which are placed at the posterior surface of the capsule and the lens fibers, which is the main substance of the crystalline lens (cortex and nucleus). The main function of the lens is to contribute to the refraction of light and its focus on the retina. When beholding a far distance object, the image of the target on the retina is well focused and we decide to focus at a near distance object, increasing the refractive power of the eye is necessary, so as to have the image of the target on the retina always well focused (accommodation). What we know today about the mechanism of accommodation are based on the theory of Helmholtz (1856) Therefore, when looking in far distance, the ciliary body is relaxed and its diameter increases, keeping the zonules fibers tense. As a result, the curvature of the crystalline lens decreases (the lens becomes flatter).

During accommodation, the ciliary muscle decreases in diameter, while the fibers get loose. Then the elastic capsule of the crystalline lens, regains its physiological curved form. Over time, the lens loses its elasticity and become harder, inflexible and more opaque in color^[7]. As a result, the lens loses the ability of accommodation. The gradual loss of the crystalline lens clarity is the pathological eye condition of cataract and is treated surgically.




Figure 1. Natural crystalline lens (left) – Crystalline lens with cataract (right)

Phacoemulsification technique

In our days, phacoemulsification^[1,2,3] is the most widespread method of surgical treatment of cataract. The main advantages of this technique are the successful removal of the crystalline lens through a small incision, the reducing of the hospitalization period, the avoidance of intraoperative and postoperative complications and the accelerating of postoperative visual rehabilitation of the patient. During this technique, the nucleus of the crystalline lens gets emulsified by a ultrasonic handpiece and the crushed parts are get removed from the anterior chamber of the eye by an irrigation-aspiration probe. Finally, a new intraocular lens (IOL), that replaces the crystalline lens, is placed inside the empty capsule.




Figure 2. Presentation of crushing the nucleus and emulsification
(<https://ljeveinstitute.org/services/cataract/phaco/>)




Figure 3. Crushing the nucleus of the crystalline lens during phacoemulsification surgery and insertion of the IOL
(http://www.kpjcfcs.com/cataract_surgery_malaysia.php)

From the April to June of 2017, at “ATTIKON” Athens University General Clinic, a clinical study took place. The purpose of the study was to examine the evolution of the refraction of eighteen patients, during the first month after phacoemulsification surgery. Also, the study aims to investigate, if the cases that were examined developed the technique’s main advantages^[4,5,6,8] Of the total of eighteen patients, eleven were women and seven were men. Their ages fluctuated between the ages of 49 and 88. They were operated by the same surgeon, who followed the same surgical process in all cases. Fifteen and thirty

days after every patient's surgical treatment, a check of the operated eye's refraction was held. Special attention was paid to the evolution of visual acuity.

Of all patients, two people showed findings of dry type of age related macular degeneration, preoperatively. In the first case (No 6), this condition affected the visual acuity of the patient and remained in low levels, postoperatively. In the second case (No 11), the visual acuity of the patient improved and didn't get affected by this pathological condition. In two people, also, appeared aspects on the Descemet's membrane fifteen days after phacoemulsification. As a result, the first patient (No 8) ended with low visual acuity and the second patient (No 18) ended with secondary myopia, because of corneal edema.

Presentation of astigmatism changes

Fifteen days after phacoemulsification, four patients developed an unimportant amount of stigmatism, between 0 - 0.25 D and six patients developed between 0.50 - 0.75 D. From the eighteen patients, six developed astigmatism between 1.00 - 1.25 D and two of them an amount of 1.75 D. In total, all cases concluded with a postoperative astigmatism. The highest rate was at 1.75 D, which is not considered important as it's going to decrease in the next few days.

At the refractive check, thirty days after the surgery, there was a little improvement in the astigmatic error. Six patients developed an unimportant rate of astigmatism between 0 - 0.25 D, five developed an astigmatic rate between 0.50 - 0.75 D and six developed astigmatism between 1.00 - 1.75 D. Finally, only one case developed a high rate of astigmatic error, at 2.00 D.

Table 1. Presentation of astigmatism changes, 15 and 30 days after phacoemulsification surgery

Number of Patients	POST-OP CYL (15 DAYS)		POST-OP CYL (30 DAYS)	
	Cyl	AXIS	Cyl	AXIS
1	-0,5	15	-0,25	80
2	-1,75	70	-1,25	75
3	-1,25	90	-1,5	106
4	-0,25	5	-0,5	160
5	0,5	170	0,5	170
6	1	75	0,75	70
7	-0,75	20	0,75	90
8	1,75	20	-1,75	100
9	0	0	0	0
10	-1	90	-1	90
11	-0,5	80	-0,25	11
12	-0,25	115	0	0
13	0	0	0,5	106
14	1,25	40	1	130

15	-1	114	-1	90
16	-0,5	45	0	0
17	-0,5	165	0	0
18	-1	96	-2	90

In conclusion, out of the eighteen clinical cases, six people developed astigmatism “with the rule”, from 0.50 to 1.50 D, four patients ended with zero astigmatism and seven patients developed “against the rule” astigmatism from 0.25 to 1.00 D. A special clinical case, was one patient (No 18), who had a final increase in astigmatism, at 2.00 D, because of a pterygium, which entering the cornea caused a refractive error.

Presentation of the spherical component of refraction changes

Fifteen days after the phacoemulsification, four patients developed a low myopic error, ranging from 0.25 to 0.50 D. Eight clinical cases, also, developed a zero sphere and the other six developed a low hyperopic error, from 0.25 to 0.50 D.

Thirty days after the surgical treatment, the noticed changes were very small, as we were already talking about a negligible sphere. As a result, the sphere remained stable in nine of the eighteen clinical cases, decreased by 0.25 D in three cases and small increase of 0.25 – 0.50 D, was noticed in six patients.

Presentation of visual acuity before and after surgery

The preoperative sphere and astigmatism weren't included as clinical data in the study, because special attention was paid to preoperative visual acuity. That's how the success of the phacoemulsification surgery gets more understandable in terms of restoring the visual capacity of the patients. Therefore, preoperatively, the visual acuity of the patients ranged from light perception (LP) to 5/10.

Table 2. Presentation of visual acuity

PATIENTS	PRE-OP VA	POST-OP VA (15 days)	POST-OP VA without glasses (30 days)	POST-OP VA with glasses (30 days)
1	0,4	0,5	0,9	1
2	0,4	0,4	0,7	0,8
3	0,4	0,6	1	1
4	0,05	0,4	0,3	1
5	0,3	0,5	0,5	0,7
6	0,15	0,7	0,7	0,7
7	0,3	0,7	0,8	0,9
8	0,05	0,4	0,6	0,9
9	0,3	0,8	0,8	1
10	0,3	0,7	0,7	0,8
11	0,2	0,7	0,9	0,9

12	0,2	0,6	0,7	0,8
13	0,4	0,6	0,7	0,9
14	0,2	0,7	1	1
15	0,5	0,9	0,9	1
16	0,05	0,8	1	1
17	0,3	0,8	0,9	0,9
18	0,3	0,4	0,6	0,8

Thirty days after the phacoemulsification surgery, twelve clinical cases developed very good to excellent visual acuity (9 – 10/10), as final postoperative visual acuity. In addition, six patients developed good visual acuity (7 – 8/10). There was complete visual acuity recovery even in patients who did not exceed 1/10 or LP, preoperatively, as the cataract was the only pathological problem of the eye. In six patients, visual acuity remained stable at 15 and 30 days and in twelve patients the vision improved. Also, thirty days after the surgery, in six out of the eighteen cases, there was a final improvement in visual acuity above two lines, when compared to visual acuity by applying the optimal refractive correction. Therefore, the use of corrective ophthalmic lenses has been proposed to them.

Table 3. Post-operative visual acuity

Number of eyes	V.A. without glasses	Number of eyes	V.A. with glasses
9	5/10→ 7/10	6	7/10→ 8/10
9	8/10→ 10/10	12	9/10→ 10/10




Chart 1. Changes of visual acuity post-operative 15 days & 30 days




Chart 2. Changes of visual acuity pre-operative and 30 days after with glasses

Statistical study

The above registration data of visual acuity, preoperatively and postoperatively, were used to perform a statistical study. Tables, statistics and paired T-tests for each case are listed below.

Sample 1	V.A. PRO OPER.
Sample 2	V.A. 15 DAYS AFTER OPER.

	Sample 1	Sample 2
Sample size	18	18
Arithmetic mean	0,2667	0,6222
95% CI for the mean	0,2006 to 0,3327	0,5430 to 0,7014
Variance	0,01765	0,02536
Standard deviation	0,1328	0,1592
Standard error of the mean	0,03131	0,03753

Paired samples t-test

Mean difference	0,3556
Standard deviation of differences	0,1901
Standard error of mean difference	0,04481
95% CI	0,2610 to 0,4501
Test statistic t	7,935
Degrees of Freedom (DF)	17
Two-tailed probability	P < 0,0001

Differences

Chi-squared test for Normal distribution of differences	accept Normality (P=0,6080) (Chi-squared=0,995 DF=2)
---	---

Table 4. Paired T-tests before phacoemulsification and 15 days after

Sample 1	V.A. 15_DAYS_AFTER_OPER.	
Sample 2	V.A. 30_DAYS_AFTER_OPER.	

	Sample 1	Sample 2
Sample size	18	18
Arithmetic mean	0,6222	0,7611
95% CI for the mean	0,5430 to 0,7014	0,6675 to 0,8548
Variance	0,02536	0,03546
Standard deviation	0,1592	0,1883
Standard error of the mean	0,03753	0,04438

Paired samples t-test

Mean difference	0,1389
Standard deviation of differences	0,1461
Standard error of mean difference	0,03443
95% CI	0,06624 to 0,2115
Test statistic t	4,034
Degrees of Freedom (DF)	17
Two-tailed probability	P = 0,0009

Differences

Chi-squared test for Normal distribution of differences	accept Normality (P=0,3231) (Chi-squared=0,976 DF=1)
---	---

Table 5. Paired T-tests 15 days and 30 days after phacoemulsification

Sample 1	V.A. PRO_OPER.	
Sample 2	V.A. 30_DAYS_AFTER_OPER.	

	Sample 1	Sample 2
Sample size	18	18
Arithmetic mean	0,2667	0,7611
95% CI for the mean	0,2006 to 0,3327	0,6675 to 0,8548
Variance	0,01765	0,03546
Standard deviation	0,1328	0,1883
Standard error of the mean	0,03131	0,04438

Paired samples t-test

Mean difference	0,4944
Standard deviation of differences	0,1955
Standard error of mean difference	0,04607
95% CI	0,3972 to 0,5916
Test statistic t	10,733
Degrees of Freedom (DF)	17
Two-tailed probability	P < 0,0001

Differences

Chi-squared test for Normal distribution of differences	accept Normality (P=0,9311) (Chi-squared=0,143 DF=2)
---	---

Table 6. Paired T-tests before phacoemulsification and 30 days after

Sample 1	V.A._30_DAYS_AFTER_OPER.	
Sample 2	V.A._30_DAYS_AFTER_OPER._WITH_GLASSES	
	Sample 1	Sample 2
Sample size	18	18
Arithmetic mean	0,7611	0,8944
95% CI for the mean	0,6675 to 0,8548	0,8419 to 0,9469
Variance	0,03546	0,01114
Standard deviation	0,1883	0,1056
Standard error of the mean	0,04438	0,02488
Paired samples t-test		
Mean difference		0,1333
Standard deviation of differences		0,1680
Standard error of mean difference		0,03961
95% CI		0,04977 to 0,2169
Test statistic t		3,367
Degrees of Freedom (DF)		17
Two-tailed probability		P = 0,0037
Differences		
Chi-squared test for Normal distribution of differences		reject Normality (P=0,0011) (Chi-squared=10,629 DF=1)

Table 7. Paired T-tests 30 days after phacoemulsification with and without applying the optimal refractive correction

Sample 1	V.A._PRO_OPER.	
Sample 2	V.A._30_DAYS_AFTER_OPER._WITH_GLASSES	
	Sample 1	Sample 2
Sample size	18	18
Arithmetic mean	0,2667	0,8944
95% CI for the mean	0,2006 to 0,3327	0,8419 to 0,9469
Variance	0,01765	0,01114
Standard deviation	0,1328	0,1056
Standard error of the mean	0,03131	0,02488
Paired samples t-test		
Mean difference		0,6278
Standard deviation of differences		0,1674
Standard error of mean difference		0,03945
95% CI		0,5446 to 0,7110
Test statistic t		15,915
Degrees of Freedom (DF)		17
Two-tailed probability		P < 0,0001
Differences		
Chi-squared test for Normal distribution of differences		accept Normality (P=0,8386) (Chi-squared=0,352 DF=2)

Table 8. Paired T-tests preoperatively and 30 days after phacoemulsification with applying the optimal refractive correction.

Conclusions

The clinical research concluded that the visual recovery of all patients was at a high level achieved without significant complications, within the first month after phacoemulsification. As a result, the phacoemulsification technique succeeded in keeping up with its promises, as better surgical treatment of ocular cataracts. This period of time was also sufficient for the eye to incorporate the IOL, heal the incisions, restore its physiology and stabilize its refraction. For this reason, the optometrist, who will be called to check the refraction of a patient who has undergone a phacoemulsification operation, must be conscious and avoid prescribing corrective ophthalmic lenses by the end of the first month.

References

1. Linebarger EJ et. al. "Phacoemulsification and modern cataract surgery", *Surv Ophthalmol*, 1999 Sep-Oct;44(2):123-47
2. Javitt JC, et. al. "Outcomes of cataract surgery. Improvement in visual acuity and subjective visual function after surgery in the first, second, and both eyes", *Arch Ophthalmol*, 1993 May;111(5):686-91
3. Charalampidou S, et al. "Prognostic indicators and outcome measures for surgical removal of symptomatic nonadvanced cataract", *Arch Ophthalmol*, 2011;129(9):1155-1161
4. Awasthi N, Guo S, Wagner BJ. "Posterior capsular opacification: a problem reduced but not yet eradicated", *Arch Ophthalmol*, 2009;127(4):555-562
5. Gothwal VK, Wright TA, Lamoureux EL, Pesudovs K, "Measuring outcomes of cataract surgery using the Visual Function Index-14", *J Cataract Refract Surg*, 2010;36(7):1181-1188
6. Lundstrom M, et. al. "The outcome of cataract surgery measured with the Catquest-9SF", *Acta Ophthalmol*, 2011;89(8):718-723
7. Chylack LT Jr, et al, "Longitudinal Study of Cataract Study Group, The Lens Opacities Classification System III", *Arch Ophthalmol*, 1993;111(6):831-836
8. Eirini Skiadaresi MD, et al, "Subjective Quality of Vision Before and After Cataract Surgery", *Arch Ophthalmol*, 2012;130(11):1377-1382
9. Rajiv B. Khandekar et al, "Visual acuity at 6 weeks after small incision cataract surgery and role of audit in predicting visual acuity", *Eur J Ophthalmol*, 2010; 20 (2): 345-352